

# RHOC Mouse mAb[5L49]

Cat NO. :A81851

#### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,ICC/IF	H,M,R	P08134	22kDa	Mouse	IgG	50ul 100ul,200ul

Applications detail:

Application

WB

1:1000-2000

ICC/IF

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human RHOC.

### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$ 

### Tissue specificity:

## Subcellular location:

Cell membrane, Lipid-anchor, Cytoplasmic side. Cleavage furrow.

#### Function:

Regulates a signal transduction pathway linking plasma membrane receptors to the assembly of focal adhesions and actin stress fibers. Serves as a microtubule-dependent signal that is required for the myosin contractile ring formation during cell cycle cytokinesis. Regulates apical junction formation in bronchial epithelial cells..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



## **Validation Data:**

## RHOC Mouse mAb[5L49] Images



Western blot (SDS PAGE) analysis of extracts from LNCaP cells.Using RHOC Mouse mAb IgG [5L49] at dilution of 1:1000 incubated at  $4^\circ\!\mathrm{C}$  over night.

View more information on http://naturebios.com