

# **HPRT Rabbit mAb [6MLX]**

Cat NO. :A85717

#### Information:

Applications	Reactivity:	UniProt ID:	ot ID: MW(kDa)	Host	Isotype	Size	
WB,IHC	H,M,R	P00492	25 kDa	Rabbit	IgG	50ul,100ul,200ul	

Applications detail:

Application

WB

1:1000-2000

IHC

The optimal dilutions should be determined by the end user

	onjug							
,	$\sim$	n		$\sim$	$\boldsymbol{\circ}$	ナヘ		
.,	.,				~		=	
_	_	• • •	_		•		-	

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human HPRT

### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$ 

### Tissue specificity:

### Subcellular location:

Cytoplasm.

#### Function:

Converts guanine to guanosine monophosphate, and hypoxanthine to inosine monophosphate. Transfers the 5-phosphoribosyl group from 5-phosphoribosylpyrophosphate onto the purine. Plays a central role in the generation of purine nucleotides through the purine salvage pathway.


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



## **Validation Data:**

## HPRT Rabbit mAb [6MLX] Images



Western blot(SDS PAGE) analysis of extracts from MCF7 CELLS .Using HPRTRabbit mAb [6MLX] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night.

View more information on http://naturebios.com