

USP9X Rabbit mAb [52RR]

Cat NO. :A87613

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	H,M,R	Q93008	270 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application	Dilution
WB	1:1000-2000
The optimal dilutions should be determined by the end user	

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of Human USP9X

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at -20°C or -80°C.

Products are valid for one natural year of receipt.Avoid repeated freeze / thaw cycles.

Tissue specificity:

Widely expressed in embryonic and adult tissues..

Subcellular location:

Cytoplasm. Cell projection, growth cone. Cytoplasm, cytoskeleton, cilium axoneme.

Function:

Deubiquitinase involved both in the processing of ubiquitin precursors and of ubiquitinated proteins (PubMed:19135894, PubMed:25944111, PubMed:18254724, PubMed:29626158). May therefore play an important regulatory role at the level of protein turnover by preventing degradation of proteins through the removal of conjugated ubiquitin (PubMed:19135894, PubMed:25944111, PubMed:18254724, PubMed:29626158). Specifically hydrolyzes 'Lys-63', 'Lys-48', 'Lys-29' and 'Lys-33'-linked polyubiquitins chains (PubMed:25944111, PubMed:18254724, PubMed:33378666). Essential component of TGF-beta/BMP signaling cascade (PubMed:19135894). Specifically deubiquitinates monoubiquitinated SMAD4, opposing the activity of E3

Introduction: **WB:** Western Blot **IP:** Immunoprecipitation **IHC:** Immunohistochemistry **ChIP:** Chromatin Immunoprecipitation **ICC/IF:** Immunocytochemistry/Immunofluorescence **F:** Flow Cytometry


Cross Reactivity: **H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Ml:** mink **C:** chicken **Dm** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Hr:** horse

For Research Use Only. Not For Use In Diagnostic Procedures.

ubiquitin-protein ligase TRIM33 (PubMed:19135894). Deubiquitinates alkylation repair enzyme ALKBH3 (PubMed:25944111). OTUD4 recruits USP7 and USP9X to stabilize ALKBH3, thereby promoting the repair of alkylated DNA lesions (PubMed:25944111). Deubiquitinates mTORC2 complex component RICTOR at 'Lys-294' by removing 'Lys-63'-linked polyubiquitin chains, stabilizing RICTOR and enhancing its binding to MTOR, thus promoting mTORC2 complex assembly (PubMed:33378666). Regulates chromosome alignment and segregation in mitosis by regulating the localization of BIRC5/survivin to mitotic centromeres (PubMed:16322459). Involved in axonal growth and neuronal cell migration (PubMed:24607389). Regulates cellular clock function by enhancing the protein stability and transcriptional activity of the core circadian protein ARNTL/BMAL1 via its deubiquitinating activity (PubMed:29626158). Deubiquitinates PEG10 (By similarity)..

Validation Data:

USP9X Rabbit mAb [52RR] Images



Western blot (SDS PAGE) analysis of extracts from 293T cells. Using USP9X Rabbit mAb [52RR] at dilution of 1:1000 incubated at 4°C over night.

View more information on <http://naturebios.com>

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.