

SEC22B Rabbit mAb [Y24L]

Cat NO. :A88828

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	H,M,R	O75396	28 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application Dilution

WB 1:1000-2000

IHC 1:1000

ICC/IF 1:100

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human SEC22B

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Subcellular location:

Endoplasmic reticulum membrane, Single-pass type IV membrane protein. Endoplasmic reticulum-Golgi intermediate compartment membrane. Golgi apparatus, cis-Golgi network membrane. Golgi apparatus,

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



SNARE involved in targeting and fusion of ER-derived transport vesicles with the Golgi complex as well as Golgi-derived retrograde transport vesicles with the ER..

Validation Data:

SEC22B Rabbit mAb [Y24L] Images



Western blot(SDS PAGE) analysis of extracts from MCF7 CELLS .Using SEC22BRabbit mAb [Y24L] at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com