

PPA2 Rabbit mAb [9818]

Cat NO. :A91165

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	H,M,R	Q9H2U2	35 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application

WB

1:1000-2000

IHC

The optimal dilutions should be determined by the end user

\sim	•					
1''	าทแ	IMATA				
\mathbf{v}	JIIIL	ıgate	•			

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human PPA2

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Detected in brain, gastric carcinoma, lung, ovary, skeletal muscle, umbilical cord blood and a cell line derived from kidney proximal tubule epithelium..

Subcellular location:

Mitochondrion.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Hydrolyzes inorganic pyrophosphate (PubMed:27523597). This activity is essential for correct regulation of mitochondrial membrane potential, and mitochondrial organization and function (PubMed:27523598)..

Validation Data:

PPA2 Rabbit mAb [9818] Images



Western blot (SDS PAGE) analysis of extracts from Human fetal kidney .Using PPA2Rabbit mAb [9818] at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com