

eICC/IF5A Rabbit mAb[YRB8]

Cat NO. :A93592

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	H,M,R	P63241	18kDa	Rabbit	IgG	50ul 100ul,200ul

Applications detail:

Application	Dilution
WB	1:1000-2000
The optimal dilutions should be determined by the end user	

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human eICC/IF5A.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at -20°C or -80°C.

Products are valid for one natural year of receipt.Avoid repeated freeze / thaw cycles.

Tissue specificity:

Expressed in umbilical vein endothelial cells and several cancer cell lines (at protein level)..

Subcellular location:

Cytoplasm. Nucleus. Endoplasmic reticulum membrane,Peripheral membrane protein,Cytoplasmic side.

Function:

mRNA-binding protein involved in translation elongation (PubMed:16987817, PubMed:17187778, PubMed:33547280). Has an important function at the level of mRNA turnover, probably acting downstream of decapping (PubMed:16987817, PubMed:17187778, PubMed:33547280). Critical for the efficient synthesis of peptide bonds between consecutive proline residues (PubMed:16987817, PubMed:17187778, PubMed:33547280). Can resolve ribosomal stalling caused by consecutive prolines during translation (PubMed:16987817, PubMed:17187778, PubMed:33547280). Involved in actin dynamics and cell cycle progression, mRNA decay and probably in a pathway involved in stress response and maintenance of cell wall

Introduction: **WB:** Western Blot **IP:** Immunoprecipitation **IHC:** Immunohistochemistry **ChIP:** Chromatin Immunoprecipitation **ICC/IF:** Immunocytochemistry/Immunofluorescence **F:** Flow Cytometry


Cross Reactivity: **H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Ml:** mink **C:** chicken **Dm** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Hr:** horse

For Research Use Only. Not For Use In Diagnostic Procedures.

integrity (PubMed:16987817, PubMed:17187778, PubMed:33547280). With syntenin SDCBP, functions as a regulator of p53/TP53 and p53/TP53-dependent apoptosis (PubMed:15371445). Regulates also TNF-alpha-mediated apoptosis (PubMed:15452064). Mediates effects of polyamines on neuronal process extension and survival (PubMed:17360499). May play an important role in brain development and function, and in skeletal muscle stem cell differentiation (By similarity). Also described as a cellular cofactor of human T-cell leukemia virus type I (HTLV-1) Rex protein and of human immunodeficiency virus type 1 (HIV-1) Rev protein, essential for mRNA export of retroviral transcripts (PubMed:8253832)..

Validation Data:

eICC/IF5A Rabbit mAb[YRB8] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cells. Using eICC/IF5A Rabbit mAb IgG [YRB8] at dilution of 1:1000 incubated at 4°C over night.

View more information on <http://naturebios.com>

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.