SCAI Rabbit mAb [ZLS3]

Cat NO. :A94620

Information:

| Applications | Reactivity: | UniProt ID: | MW(kDa) | Host | Isotype | Size |
|---------------|-------------|-------------|---------|--------|---------|-------------|
| WB,IHC,ICC/IF | H,M,R | Q8N9R8 | 66 kDa | Rabbit | lgG | 100ul,200ul |

Applications detail:

| Application | Dilution | | |
|--|-------------|--|--|
| WB | 1:1000-2000 | | |
| ІНС | 1:100 | | |
| ICC/IF | 1:100 | | |
| The optimal dilutions should be determined by the end user | | | |

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human SCAI

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Membrane, Single-pass membrane protein. Nucleus. Cytoplasm.

Function:

Tumor suppressor which functions to suppress MRTFA-induced SRF transcriptional activity. May function in the RHOA-DIAPH1 signal transduction pathway and regulate cell migration through transcriptional regulation of ITGB1..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Validation Data:

SCAI Rabbit mAb [ZLS3] Images



Western blot (SDS PAGE) analysis of extracts from Human thymus.Using SCAIRabbit mAb [ZLS3] at dilution of 1:1000 incubated at 4° over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.