

RAGE Rabbit mAb [3EUJ]

Cat NO. :A95147

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	H,M,R	Q9UQ07	45,55 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

ApplicationDilutionWB1:1000-2000IHC1:100ICC/IF1:100The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human RAGE

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Expressed in heart, brain, lung, kidney, and pancreas, and at very low levels in placenta, liver and skeletal muscle.

Detected in retina.

Subcellular location:

Cytoplasm. Cell projection, cilium. Nucleus.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Able to phosphorylate several exogenous substrates and to undergo autophosphorylation. Negatively regulates cilium length in a cAMP and mTORC1 signaling-dependent manner..

Validation Data:

RAGE Rabbit mAb [3EUJ] Images



Western blot (SDS PAGE) analysis of extracts from Mouse lung lysates with 5% NFDM/TBST.Using RAGERabbit mAb [3EUJ] at dilution of 1:1000 incubated

View more information on http://naturebios.com