

# Collagen Type I Mouse mAb[4L77]

Cat NO. :A96544

### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	н	P02452	130kda	Mouse	IgG	50ul 100ul,200ul

Applications detail:

Application	Dilution			
WB	1:1000-2000			
IHC	1:100			
ICC/IF	1:100			
The optimal dilutions should be determined by the end user				

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

**Endogenous** 

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human Collagen Type I.

### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

### Tissue specificity:

Forms the fibrils of tendon, ligaments and bones. In bones the fibrils are mineralized with calcium hydroxyapatite.

## Subcellular location:

Secreted, extracellular space, extracellular matrix.

### Function:

Type I collagen is a member of group I collagen (fibrillar forming collagen).


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



## **Validation Data:**

## Collagen Type I Mouse mAb[4L77] Images



Western blot (SDS PAGE) analysis of extracts from human cervical cancer tissue. Using Collagen Type I Mouse mAb IgG [4L77] at dilution of 1:1000 incubated at

View more information on http://naturebios.com